

**Name:**  
**Room:**  
**Age:**  
**Sex:** M F    **Pronouns:**

**CC:**  
**HPI:**

**Psych ROS**

Mood (depressed, hopeless, worthless, etc.):  
 Depressed (1-10):  
 Anxious (1-10):  
 Panic Attack/Sympt:  
 Sleep: ↑ ↓ ↔    Hrs:  
 Interest: ↑ ↓ ↔    **Guilt:** Y N  
 Energy: ↑ ↓ ↔    **Concentration:** ↑ ↓ ↔  
 Memory: ↑ ↓ ↔    **Appetite:** ↑ ↓ ↔  
 Weight: ↑ ↓ ↔ (Amt/time):  
 Psychomotor Agitation/Restless: Y N  
 Manic (Distractibility, Indiscreet (risky behavior),  
 Grandiosity, Flight of ideas, Active w/ goal, Sleep deficit, Talk  
 (pressured))  
 OCD: Y N  
 Suicidal: Y N plan:  
 Homicidal: Y N plan:  
 AH: Y N  
 VH: Y N  
 Paranoia: Y N  
 Delusion: Y N

**Past Med Hx**

HTN HLD DM Thyroid Renal Liver  
 Concussions/TBI:  
 Other Dx:

**Current Meds:**

Med	Dose

Allergies:

**Past Psych Hx**

Prior Dx:  
 Past Hosp:  
 Outpatient Services:  
 Therapist:  
 Psychiatrist:  
 Previous Suicides/Methods/Reasons:  
 - Total Attempts #:  
 - Most Common Method:  
 - First Attempt  
 - Last Attempt:  
 - Self-Harm Behaviors:  
 Trauma (sexual, emotional, physical):  
 Flashbacks/Nightmares: Y N

**Past Tried Meds**

Med	Dose	D/C reason

**Substance Use Hx:**

- EtOH (type/amt):  
 - Tobacco (ppd/yr):  
 - THC (Smoke, oil, edible):  
 - Cocaine, LSD, Heroin, Benzo, Meth, PCP,  
 K2, other: \_\_\_\_\_  
 Detox: Y N  
 Rehab: Y N  
 Seizure Hx: Y N

**Fam Hx**

Fam Psych Dx Hx:  
 Fam Suicide Hx:  
 Fam Drug Abuse Hx:

**Social Hx**

Highest Education:  
 Occupation:  
 Legal/Probation:  
 Military: Y N  
 Firearms: Y N locked/ammo separated  
 Housing:  
 Relationship Status:  
 Children:  
 Stressors:

Support:

**Additional Notes**